

FORM PTO-1390 (Modified)  
(REV 11-2000)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

**TRANSMITTAL LETTER TO THE UNITED STATES**  
**DESIGNATED/ELECTED OFFICE (DO/EO/US)**  
**CONCERNING A FILING UNDER 35 U.S.C. 371**

209861US0PCT

U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR

09/868338

INTERNATIONAL APPLICATION NO.

PCT/JP99/07079

INTERNATIONAL FILING DATE

16 December 1999

PRIORITY DATE CLAIMED


18 December 1998

TITLE OF INVENTION

ABC TRANSPORTER AND GENE CODING FOR THE SAME

APPLICANT(S) FOR DO/EO/US

Sohei KANNO, et al



Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), (9) and (24) indicated below.
4. ☒ The US has been elected by the expiration of 19 months from the priority date (Article 31).
5. ☒ A copy of the International Application as filed (35 U.S.C. 371 (c) (2))
  - a. ☐ is attached hereto (required only if not communicated by the International Bureau).
  - b. ☒ has been communicated by the International Bureau.
  - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
- ☒ An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)).
  - a. ☒ is attached hereto.
  - b. ☐ has been previously submitted under 35 U.S.C. 154(d)(4).
- ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371 (c)(3))
  - a. ☐ are attached hereto (required only if not communicated by the International Bureau).
  - b. ☐ have been communicated by the International Bureau.
  - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
  - d. ☒ have not been made and will not be made.
- ☐ An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
- ☒ An oath or declaration of the inventor(s) (35 U.S.C. 371 (c)(4)).
- ☒ An English language translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371 (c)(5)).
- ☐ A copy of the International Preliminary Examination Report (PCT/IPEA/409).
- ☒ A copy of the International Search Report (PCT/ISA/210).

**Items 13 to 20 below concern document(s) or information included:**

13. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
14. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
15. ☐ A **FIRST** preliminary amendment.
16. ☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
17. ☐ A substitute specification.
18. ☐ A change of power of attorney and/or address letter.
19. ☐ A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821 - 1.825.
20. ☐ A second copy of the published international application under 35 U.S.C. 154(d)(4).
21. ☐ A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4).
22. ☐ Certificate of Mailing by Express Mail
23. ☒ Other items or information:

Request for Consideration of Documents Cited in International Search Report

Notice of Priority

PCT/IB/304, PCT/IB/308

Sequence Listing (10 Pages)

U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR 1.492(a)(1) - (5)) : <b>097868338</b>		INTERNATIONAL APPLICATION NO. <b>PCT/JP99/07079</b>		ATTORNEY'S DOCKET NUMBER <b>209861US0PCT</b>	
---	--	--	--	---	--

24. The following fees are submitted:.				<b>CALCULATIONS PTO USE ONLY</b>	
<b>BASIC NATIONAL FEE ( 37 CFR 1.492 (a) (1) - (5) ) :</b> <input type="checkbox"/> Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO ..... <b>\$1000.00</b> <input checked="" type="checkbox"/> International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO ..... <b>\$860.00</b> <input type="checkbox"/> International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO ..... <b>\$710.00</b> <input type="checkbox"/> International preliminary examination fee (37 CFR 1.482) paid to USPTO but all claims did not satisfy provisions of PCT Article 33(1)-(4) ..... <b>\$690.00</b> <input type="checkbox"/> International preliminary examination fee (37 CFR 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4) ..... <b>\$100.00</b> <div style="text-align: right;"><b>ENTER APPROPRIATE BASIC FEE AMOUNT =</b></div>				<b>\$860.00</b>	
Surcharge of <b>\$130.00</b> for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492 (e)).					
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE		
Total claims	14 - 20 =	0	x \$18.00	<b>\$0.00</b>	
Independent claims	7 - 3 =	4	x \$80.00	<b>\$320.00</b>	
Multiple Dependent Claims (check if applicable). <input type="checkbox"/>				<b>\$0.00</b>	
<b>TOTAL OF ABOVE CALCULATIONS =</b>				<b>\$1,180.00</b>	
<input type="checkbox"/> Applicant claims small entity status. (See 37 CFR 1.27). The fees indicated above are reduced by 1/2.				<b>\$0.00</b>	
<b>SUBTOTAL =</b>				<b>\$1,180.00</b>	
Processing fee of <b>\$130.00</b> for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492 (f)).				<b>\$0.00</b>	
<b>TOTAL NATIONAL FEE =</b>				<b>\$1,180.00</b>	
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) (check if applicable). <input type="checkbox"/>				<b>\$0.00</b>	
<b>TOTAL FEES ENCLOSED =</b>				<b>\$1,180.00</b>	
				<b>Amount to be: refunded</b>	\$
				<b>charged</b>	\$

a. ☒ A check in the amount of **\$1,180.00** to cover the above fees is enclosed.

b. ☐ Please charge my Deposit Account No. \_\_\_\_\_ in the amount of \_\_\_\_\_ to cover the above fees. A duplicate copy of this sheet is enclosed.

c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. **15-0030** A duplicate copy of this sheet is enclosed.

d. ☐ Fees are to be charged to a credit card. **WARNING:** Information on this form may become public. **Credit card information should not be included on this form.** Provide credit card information and authorization on PTO-2038.

**NOTE:** Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

Surinder Sachar  
Registration No. 34,423

*Surinder Sachar*

SIGNATURE

**Norman F. Oblon**

NAME

**24,618**

REGISTRATION NUMBER

**6-18-01**

DATE

DOCKET NO.: 209861US-0PCT

**IN THE UNITED STATES PATENT & TRADEMARK OFFICE**

IN RE APPLICATION OF :  
Sohei KANNNO et al : ATTN: BOX SEQUENCE  
SERIAL NO: 09/868,338 :  
FILED: June 18, 2001  
FOR: ABC TRANSPORTER AND GENE  
CODING FOR THE SAME

**PRELIMINARY AMENDMENT AND STATEMENT**

ASSISTANT COMMISSIONER FOR PATENTS  
WASHINGTON, D.C. 20231

SIR:

Responsive to the Notice to Comply dated July 23, 2001, Applicants submit herewith  
amendments to the specification, a substitute Sequence Listing, and a corresponding  
Computer-Readable Sequence Listing.

**IN THE SPECIFICATION**

Please replace the paragraph beginning on page 4, line 25, through page 5, line 11,  
with the following paragraph:

--PCR (polymerase chain reaction) is performed by using chromosomal DNA of  
*Brevibacterium lactofermentum* such as *Brevibacterium lactofermentum* ATCC13869 as a  
template and primers having nucleotide sequences of regions in the *gltBD* genes of  
*Escherichia Coli* K-12 (*Gene*, vol. 60, pp. 1-11 (1987) and yeast (*Saccharomyces cerevisiae*,  
GenBank Accession No. X89221) exhibiting high homology, for example, those having  
nucleotide sequences of SEQ ID NOS: 1 and 2 shown in Sequence Listing, to obtain a DNA

fragment of about 1.4 kb. *Brevibacterium lactofermentum* ATCC13869 can be obtained from ATCC (the American Type Culture Collection: 10801 University Boulevard, Manassas, VA 20110-2209, United States of America).--

Please replace the paragraph on page 9, line 24, through page 10, line 26, with the following paragraph:

--A DNA coding for substantially the same protein as a constituent of ABC transporter can be obtained by expressing DNA having such a mutation as described above in an appropriate cell, and examining characteristics of an expressed product. A DNA coding for substantially the same protein as a constituent of ABC transporter can also be obtained by isolating a DNA hybridizable with a nucleotide sequence coding for each constituent or a probe prepared from such a nucleotide sequence, for example, the nucleotide sequence of nucleotide numbers 1117 to 1725 in SEQ ID NO: 7 or a probe prepared from this nucleotide sequence, for ATPase under a stringent condition, and coding for a protein having the characteristics of the constituent from a DNA coding for each protein having mutation or from a cell harboring it. The "stringent condition" referred to herein is a condition under which a so-called specific hybrid is formed, but a non-specific hybrid is not formed. It is difficult to clearly define this condition by using numerical values. However, for example, the stringent condition includes a condition under which two of DNAs having high homology, for example, two of DNAs having homology of not less than 60% are hybridized with each other, but two of DNAs having homology lower than the above level are not hybridized with each other. Alternatively, the stringent condition is exemplified by a hybridization condition represented by salt concentrations of 1 x SSC, 0.1% SDS, preferably

0.1 x SSC, 0.1% SDS, at 60°C, which is an ordinary condition of washing in Southern hybridization.--

Page 26 (Abstract), after the last line, beginning on the next page, please replace page 27 to page 36 of the original Sequence Listing with the substitute Sequence Listing attached herewith.

#### **IN THE CLAIMS**

Please amend the claims as follows:

--4. (Amended) The DNA according to Claim 3, wherein the stringent condition is a condition in which hybridization is performed at 60°C and a salt concentration corresponding to 1 x SSC and 0.1% SDS.

8. (Amended) The DNA according to Claim 7, wherein the stringent condition is a condition in which hybridization is performed at 60°C and a salt concentration corresponding to 1 x SSC and 0.1% SDS.

12. (Amended) The DNA according to Claim 11, wherein the stringent condition is a condition in which hybridization is performed at 60°C and at a salt concentration corresponding to 1 x SSC and 0.1% SDS.

#### **REMARKS**

Claims 1-14 are active in the present application. Claims 4, 8 and 12 have been amended. Support for the amendment is found, for example, on page 10, lines 12-26. The specification has been amended to correct typographical or clerical errors.

Applicants have now submitted a substitute Sequence Listing and a corresponding Computer-Readable Sequence Listing. Contents of the paper copy of the substitute Sequence Listing and the Computer-Readable Sequence Listing are identical. Support for all the sequences listed in the substitute Sequence Listing can be found in the present application. No new matter is introduced by the submission of the substitute Sequence Listing and the Computer-Readable Sequence Listing.

Applicants submit that this application is now in condition for examination on the merits. Early notice to this effect is earnestly solicited.

Respectfully submitted,

OBLON, SPIVAK, McCLELLAND,  
MAIER & NEUSTADT, P.C.



Norman F. Oblon  
Attorney of Record  
Registration No. 24,618

Daniel J. Pereira, Ph.D.  
Registration No. 45,518



**22850**

TEL: (703) 413-3000  
FAX: (703) 413-2220

NFO:DJP:TWB:smn  
I:\atty\Twb\209861.pr.wpd

**MARKED-UP COPY OF**  
**PRELIMINARY AMENDMENT AND STATEMENT**

**IN THE SPECIFICATION**

Please replace the paragraph beginning on page 4, line 25, through page 5, line 11, with the following paragraph:

--PCR (polymerase chain reaction) is performed by using chromosomal DNA of *Brevibacterium lactofermentum* such as *Brevibacterium lactofermentum* ATCC13869 as a template and primers having nucleotide sequences of regions in the *gltBD* genes of *Escherichia Coli* K-12 (*Gene*, vol. 60, pp. 1-11 (1987) and yeast (*Saccharomyces cerevisiae*, GenBank Accession No. X89221) exhibiting high homology, for example, those having nucleotide sequences of SEQ ID NOS: 1 and 2 shown in Sequence Listing, to obtain a DNA fragment of about 1.4 kb. *Brevibacterium lactofermentum* ATCC13869 can be obtained from ATCC (the American Type Culture Collection: [12301 Parklawn Drive, Rockville, Maryland, 20852] 10801 University Boulevard, Manassas, VA 20110-2209, United States of America).--

Please replace the paragraph on page 9, line 24, through page 10, line 26, with the following paragraph:

--A DNA coding for substantially the same protein as a constituent of ABC transporter can be obtained by expressing DNA having such a mutation as described above in an appropriate cell, and examining characteristics of an expressed product. A DNA coding for substantially the same protein as a constituent of ABC transporter can also be obtained by

isolating a DNA hybridizable with a nucleotide sequence coding for each constituent or a probe prepared from such a nucleotide sequence, for example, the nucleotide sequence of nucleotide numbers 1117 to 1725 in SEQ ID NO: 7 or a probe prepared from this nucleotide sequence, for ATPase under a stringent condition, and coding for a protein having the characteristics of the constituent from a DNA coding for each protein having mutation or from a cell harboring it. The "stringent condition" referred to herein is a condition under which a so-called specific hybrid is formed, but a non-specific hybrid is not formed. It is difficult to clearly define this condition by using numerical values. However, for example, the stringent condition includes a condition under which two of DNAs having high homology, for example, two of DNAs having homology of not less than [40%] 60% are hybridized with each other, but two of DNAs having homology lower than the above level are not hybridized with each other. Alternatively, the stringent condition is exemplified by a hybridization condition represented by salt concentrations of 1 x SSC, 0.1% SDS, preferably 0.1 x SSC, 0.1% SDS, at 60°C, which is an ordinary condition of washing in Southern hybridization.--

#### **IN THE CLAIMS**

--4. (Amended) The DNA according to Claim 3, wherein the stringent condition is a condition in which [washing] hybridization is performed at 60°C and a salt concentration corresponding to 1 x SSC and 0.1% SDS.

8. (Amended) The DNA according to Claim 7, wherein the stringent condition is a condition in which [washing] hybridization is performed at 60°C and a salt concentration corresponding to 1 x SSC and 0.1% SDS.



12. (Amended) The DNA according to Claim 11, wherein the stringent condition is a condition in which [washing] hybridization is performed at 60°C and at a salt concentration corresponding to 1 x SSC and 0.1% SDS.--

## SPECIFICATION

## ABC TRANSPORTER AND GENE CODING FOR THE SAME

5                    Technical Field

The present invention relates to a novel ABC transporter and a gene coding for a protein that is a constituent of the ABC transporter. The gene can be  
10 utilized for breeding of a microorganism showing modified transport of amino acids across a cell membrane and so forth.

15                    Background Art

There are several mechanisms are known for transport of substances such as an amino acids or ions through cell membranes. As one of such mechanisms, the ATP-binding cassette superfamily (ABC transporters) is  
20 known (C.F. Higgins, *Ann. Rev. Cell Biol.*, 8, 67 (1992)).

The ATP-binding cassettes constitute a group of proteins having an ATP-binding domain including a transmembrane domain. Their physiological function is primarily uptake of substances into a cell, but the ATP-  
25 binding cassette is considered to also participate in excretion of substances to some extent. In bacteria, they usually contain, as constituents, membrane proteins

09868338-061801

(membrane components), proteins that are present inside the membrane and have the ATPase activity, and binding proteins that are present outside the membrane and bound to substances. The membrane proteins and proteins  
5 having the ATPase activity form a polymer complex. It is said that the substance excretion system lacks a binding protein bound to a substance to be transported (Reizer, J. et al., *Prot. Sci.* 1, 1326 (1992)).

Since the ABC transporters or constituents thereof  
10 are involved in transport of substances, it is considered that characteristics of a cell concerning substance transport can be modified by modifying expression of genes coding for them.

Structures of various ABC transporter genes in  
15 bacteria such as *Escherichia coli* have been analyzed, and it is known that each gene coding for constituent of an ABC transporter forms an operon. In coryneform bacteria, however, most of genes coding for ABC transporters or constituents thereof involved in  
20 transport of amino acids across membranes remain unknown.

#### Disclosure of the Invention

The inventors of the present invention cloned a  
25 gene coding for an enzyme involved in one of L-glutamic acid biosynthetic pathways, glutamine-oxoglutarate aminotransferase (also referred to as glutamate synthase,

09060338-061001

abbreviated as "GOGAT" hereinafter) for the purpose of breeding of coryneform bacteria producing L-glutamic acid. In this process, the inventors accidentally found that a DNA fragment containing a gene coding for GOGAT  
5 (gltBD) contained a gene coding for an ABC transporter considered to be involved in transport of amino acids, and thus accomplished the present invention.

That is, the present invention provides a protein, which is a constituent of ABC transporter, and a DNA  
10 coding for it.

A first constituent of ABC transporter according to the present invention is a protein defined in the following (A) or (B):

(A) a protein which has the amino acid sequence of  
15 SEQ ID NO: 8 shown in Sequence Listing;

(B) a protein which has the amino acid sequence of SEQ ID NO: 8 shown in Sequence Listing including substitution, deletion, insertion, addition or inversion of one or several amino acids, and constitutes an ABC  
20 transporter.

A second constituent of ABC transporter according to the present invention is a protein defined in the following (C) or (D):

(C) a protein which has the amino acid sequence of  
25 SEQ ID NO: 9 shown in Sequence Listing;

(D) a protein which has the amino acid sequence of SEQ ID NO: 9 shown in Sequence Listing including

substitution, deletion, insertion, addition or inversion of one or several amino acids, and has ATPase activity of ABC transporter.

A third constituent of ABC transporter according to the present invention is a protein defined in the following (E) or (F):

(E) a protein which has the amino acid sequence of SEQ ID NO: 10 shown in Sequence Listing;

(F) a protein which has the amino acid sequence of SEQ ID NO: 10 shown in Sequence Listing including substitution, deletion, insertion, addition or inversion of one or several amino acids, and constitutes an ABC transporter.

The present invention also provides DNAs coding for the aforementioned proteins that are constituents of ABC transporter.

The present invention further provides an operon coding for an ABC transporter.

Hereafter, the present invention will be explained in detail.

The DNA of the present invention was found from *Brevibacterium lactofermentum* as an ORF present in the neighborhood of the *gltBD* gene and can be obtained as follows.

PCR (polymerase chain reaction) is performed by using chromosomal DNA of *Brevibacterium lactofermentum* such as *Brevibacterium lactofermentum* ATCC13869 as a

09060333, 061304

template and primers having nucleotide sequences of regions in the *gltBD* genes of *Escherichia Coli* K-12 (Gene, vol. 60, pp.1-11 (1987) and yeast (*Saccharomyces cerevisiae*, GenBank Accession No. X89221) exhibiting  
5 high homology, for example, those having nucleotide sequences of SEQ ID NOS: 1 and 2 shown in Sequence Listing, to obtain a DNA fragment of about 1.4 kb. *Brevibacterium lactofermentum* ATCC13869 can be obtained from ATCC (the American Type Culture Collection: 12301  
10 Parklawn Drive, Rockville, Maryland 20852, United States of America).

Subsequently, colony hybridization of a chromosomal DNA library of *Brevibacterium lactofermentum* ATCC13869 is performed by using the PCR-amplified  
15 fragment obtained as described above as a probe to obtain a DNA fragment hybridizable with the probe. Thus, the DNA of the present invention can be obtained together with the *gltBD* gene. If chromosomal DNA digested with *HindIII* is used in the preparation of the  
20 chromosomal DNA library, the DNA fragment can be obtained as a fragment of about 14 kb in length.

The above DNA fragment contains the *gltBD* gene and two open reading frames (ORFs) downstream the *gltBD* gene in the inverted direction with respect to the *gltBD* gene  
25 from the end. These ORFs correspond to the second ORF and third ORF, respectively, among the ORFs included in the nucleotide sequence of SEQ ID NO: 7.

As shown in examples described later, it is possible that the aforementioned two ORFs form an operon together with another ORF that exists upstream from them. This ORF corresponds to the first ORF among the ORFs included in the nucleotide sequence of SEQ ID NO: 7. This first ORF can be obtained as a DNA fragment of about 1.8 kb by PCR using chromosomal DNA of *Brevibacterium lactofermentum*, for example, the *Brevibacterium lactofermentum* ATCC13869, as a template and nucleotide sequences of SEQ ID NOS: 5 and 6 shown in Sequence Listing as primers. In this DNA fragment, a region estimated to be a promoter region exists in the upstream of the target ORF.

The nucleotide sequence shown in SEQ ID NO: 7 is obtained by ligating a nucleotide sequence (1.3 kb) in the aforementioned DNA fragment of about 14 kb with a nucleotide sequence (1.1 kb) in the aforementioned DNA fragment of about 1.8 kb.

Since the nucleotide sequences of the above ORFs and nucleotide sequences of flanking regions have been revealed, the above ORFs can also be obtained by PCR using oligonucleotides prepared based on such nucleotide sequences as primers.

Usual methods well known to those skilled in the art can be employed for preparation of chromosomal DNA, construction of chromosomal DNA library, hybridization, PCR, preparation of plasmid DNA, digestion and ligation

of DNA, transformation, design of oligonucleotides to be  
used as primers and so forth. These methods are  
described in Sambrook, J., Fritsch, E.F., Maniatis, T.,  
Molecular Cloning, Cold Spring Harbor Laboratory Press,  
5 1.21 (1989) and so forth.

The aforementioned second ORF and amino acid  
sequence encoded thereby were compared with known  
sequences for homology. The used databases were EMBL  
and SWISS-PROT. As a result, these sequences exhibited  
10 homology to already reported ATPase proteins  
constituting ABC transporters responsible for transport  
of the amino acids listed in Table 1 and genes coding  
for them. It is possible that the three ORFs containing  
these sequences form an operon.

15

0966333.061801



Table 1

Gene	Substance to be transported	Origin	Reference	Homology
artP	Arginine	<i>E. coli</i>	J.Bacteriol.175: 3687-3688 (1993)	31.0%
artP	Arginine	<i>Haemophilus Influenzae</i>	Science 269: 496-512 (1995)	31.8%
glnQ	Glutamine	<i>Bacillus Stearothermophilus</i>	J.Bacteriol.173: 4877-4888 (1991)	35.4%
glnQ	Glutamine	<i>E. coli</i>	Mol.Gen.Genet.205: 260-269 (1986)	33.5%
GltL	Glutamic acid/Aspartic acid	<i>E. coli</i>	GeneBank Accession No.U10981	33.5%
gltL	Glutamic acid/Aspartic acid	<i>Haemophilus influenzae</i>	Science 269: 496-512 (1995)	31.2%
gluA	Glutamic acid	<i>Corynebacterium glutamicum</i>	J.Bacteriol.177: 1152-1158	34.4%
hisP	Histidine	<i>E. coli</i>	Nature 298: 723-727 (1982)	33.0%
hisP	Histidine	<i>Salmonella typhimurium</i>	Nucleic acids Res.15: 8568-8568	34.4%

The gene coding for a constituent of ABC

5 transporter according to the present invention may be one coding for an ATP-binding protein including substitution, deletion, insertion, addition or inversion of one or several amino acids at one or a plurality of positions so long as characteristics of the encoded

10 protein are not deteriorated. The number meant by the term "several" used herein may vary depending on locations of amino acid residues in the three-dimensional structure of proteins and kinds of amino acid residues. This is due to the fact that there are

15 highly analogous amino acids among amino acids such as isoleucine and valine, and difference among such amino

0986338.061301

acids does not substantially affect the three-dimensional structure of proteins.

Such a DNA encoding a protein substantially the same as a constituent of ABC transporter as mentioned above can be obtained by modifying a nucleotide sequence by, for example, site-directed mutagenesis so that the amino acid residues of a specific site should include substitution, deletion, insertion, addition or inversion. Such a modified DNA as mentioned above can also be obtained by an already known mutagenesis treatment. Examples of the mutagenesis treatment include *in vitro* treatment of DNA coding for each protein with hydroxylamine etc., treatment of a microorganism having DNA coding for each protein, for example, genus *Escherichia*, by ultraviolet irradiation or with a mutagenizing agent used for usual mutagenesis treatment such as N-methyl-N'-nitro-N-nitrosoguanidine (NTG) or nitrous acid.

The substitution, deletion, insertion, addition or inversion of nucleotides described above also includes mutations (mutant or variant) that naturally occurring due to individual difference, difference in species or genera of a microorganism having each constituent.

A DNA coding for substantially the same protein as a constituent of ABC transporter can be obtained by expressing DNA having such a mutation as described above in an appropriate cell, and examining characteristics of

09060338 "061301

an expressed product. A DNA coding for substantially the same protein as a constituent of ABC transporter can also be obtained by isolating a DNA hybridizable with a nucleotide sequence coding for each constituent or a probe prepared from such a nucleotide sequence, for example, the nucleotide sequence of nucleotide numbers 1117 to 1725 in SEQ ID NO: 7 or a probe prepared from this nucleotide sequence, for ATPase under a stringent condition, and coding for a protein having the characteristics of the constituent from a DNA coding for each protein having mutation or from a cell harboring it. The "stringent condition" referred to herein is a condition under which a so-called specific hybrid is formed, but a non-specific hybrid is not formed. It is difficult to clearly define this condition by using numerical values. However, for example, the stringent condition includes a condition under which two of DNAs having high homology, for example, two of DNAs having homology of not less than 40% are hybridized with each other, but two of DNAs having homology lower than the above level are not hybridized with each other. Alternatively, the stringent condition is exemplified by a hybridization condition represented by salt concentrations of 1 x SSC, 0.1% SDS, preferably 0.1 x SSC, 0.1% SDS, at 60°C, which is an ordinary condition of washing in Southern hybridization.

Those genes hybridizable under the condition as

described above include those having a stop codon generated in the genes, and those having no activity due to mutation of the active center. However, such mutant genes can be easily removed by using a commercially available activity expression vector to examine the characteristics of the expressed product.

The DNA coding for a constituent of ABC transporter according to the present invention and an operon of ABC transporter (hereafter, these may be referred to simply as "gene of the present invention") can be utilized in breeding of coryneform bacteria. That is, since the ABC transporter of the present invention or a constituent thereof is considered to be involved in transport of amino acids, characteristics of a cell concerning transport of amino acids can be modified by modifying expression of these genes.

Coryneform bacteria to which the present invention is applicable include those bacteria having been hitherto classified into the genus *Brevibacterium* but united into the genus *Corynebacterium* at present (*Int. J. Syst. Bacteriol.*, 41, 255 (1981)), and include bacteria belonging to the genus *Brevibacterium* closely relative to the genus *Corynebacterium*. Examples of such coryneform bacteria are mentioned below.

*Corynebacterium acetoacidophilum*  
*Corynebacterium acetoglutamicum*  
*Corynebacterium alkanolyticum*

*Corynebacterium callunae*

*Corynebacterium glutamicum*

*Corynebacterium lilium* (*Corynebacterium glutamicum*)

5 *Corynebacterium melassecola*

*Corynebacterium thermoaminogenes*

*Corynebacterium herculis*

*Brevibacterium divaricatum* (*Corynebacterium glutamicum*)

10 *Brevibacterium flavum* (*Corynebacterium glutamicum*)

*Brevibacterium immariophilum*

*Brevibacterium lactofermentum* (*Corynebacterium glutamicum*)

*Brevibacterium roseum*

15 *Brevibacterium saccharolyticum*

*Brevibacterium thiogenitalis*

*Brevibacterium album*

*Brevibacterium cerium*

*Microbacterium ammoniophilum*

20 Specifically, the following strains can be exemplified.

*Corynebacterium acetoacidophilum* ATCC 13870

*Corynebacterium acetoglutamicum* ATCC 15806

*Corynebacterium alkanolyticum* ATCC21511

25 *Corynebacterium callunae* ATCC 15991

*Corynebacterium glutamicum* ATCC 13020, 13032,

13060

0986838-064801

*Corynebacterium lilium* (*Corynebacterium glutamicum*) ATCC 15990

*Corynebacterium melassecola* ATCC 17965

*Corynebacterium thermoaminogenes* AJ12340 (FERM BP-  
5 1539)

*Corynebacterium herculis* ATCC13868

*Brevibacterium divaricatum* (*Corynebacterium glutamicum*) ATCC 14020

*Brevibacterium flavum* (*Corynebacterium glutamicum*)  
10 ATCC 13826, ATCC 14067

*Brevibacterium immariophilum* ATCC 14068

*Brevibacterium lactofermentum* (*Corynebacterium glutamicum*) ATCC 13665, ATCC 13869

*Brevibacterium roseum* ATCC 13825

15 *Brevibacterium saccharolyticum* ATCC 14066

*Brevibacterium thiogenitalis* ATCC 19240

*Brevibacterium album* ATCC15111

*Brevibacterium cerium* ATCC15112

*Microbacterium ammoniaphilum* ATCC15354

20 Methods of modifying a gene coding for an ABC  
transporter or a constituent thereof include  
amplification or disruption of the gene. The gene or  
the like can be amplified by transforming a coryneform  
bacterium with a recombinant vector obtained by ligating  
25 the gene to a vector such as a plasmid. At this time,  
amplification efficiency can be improved by using a  
multiple copy type vector. Examples of such a vector

0986338-064301

include plasmids autonomously replicable in coryneform bacterium including those mentioned below.

pAM330 (refer to Japanese Patent Laid-Open (Kokai) No. 58-67699)

5 pHM1519 (refer to Japanese Patent Laid-Open No. 58-77895)

pAJ655 (refer to Japanese Patent Laid-Open No. 58-192900)

10 pAJ611 (refer to Japanese Patent Laid-Open No. 58-192900)

pAJ1844 (refer to Japanese Patent Laid-Open No. 58-192900)

pCG1 (refer to Japanese Patent Laid-Open No. 57-134500)

15 pCG2 (refer to Japanese Patent Laid-Open No. 58-35197)

pCG4 (refer to Japanese Patent Laid-Open No. 57-183799)

20 pCG11 (refer to Japanese Patent Laid-Open No. 57-183799)

Coryneform bacteria can be transformed by the electric pulse method (refer to Japanese Patent Laid-Open No. 2-207791).

25 The gene can also be amplified by allowing multiple copies of the gene of the present invention to exist on chromosomal DNA of a host such as those mentioned above. Multiple copies of a target gene can

09868338-064804

be introduced into chromosomal DNA of coryneform  
bacterium by homologous recombination utilizing multiple  
copies of sequences existing on chromosomal DNA as  
targets (Experiments in Molecular Genetics, Cold Spring  
5 Harbor Laboratory Press (1972); Matsuyama, S. and  
Mizushima, S., *J. Bacteriol.*, 162, 1196 (1985)). As  
sequences of which multiple copies exist on the  
chromosomal DNA, repetitive DNA and inverted repeats  
that exist at an end of transposable element can be used.  
10 As disclosed in Japanese Patent Laid-open No. 2-109985,  
it is also possible to insert the target gene into  
transposon, and allow it to transfer to introduce  
multiple copies thereof into the chromosomal DNA.

Further, expression of the gene can be modified by  
15 replacing an expression regulatory sequence of the gene  
originally present on a chromosome, such as a promoter,  
with a stronger one or one having weak functions.

Moreover, gene disruption methods by homologous  
recombination have already been established, and the  
20 gene can be disrupted by a method using linear DNA or a  
temperature sensitive plasmid.

#### Best Mode for Carrying out the Invention

25 Hereafter, the present invention will be explained  
in more detail with reference to the following examples.

09868338-051801



(1) Cloning of *gltBD* gene of *Brevibacterium lactofermentum* ATCC13869

09668333 061804

A region of *gltB* gene products of *Escherichia coli* and yeast showing high homology for amino acid sequence was selected, and a nucleotide sequence was deduced from the sequence, oligonucleotides shown as SEQ ID NOS: 1 and 2 were synthesized. Separately, chromosomal DNA of *Brevibacterium lactofermentum* ATCC13869 was prepared by using a Bacterial Genomic DNA Purification Kit (produced by Advanced Genetic Technologies Corp.). PCR was performed by using this chromosomal DNA as a template and the oligonucleotides as primers under the standard reaction conditions described in "PCR Technology", p. 8, Ed. by Henry Ehrlich, Stockton Press, 1989. The PCR product was subjected to agarose gel electrophoresis, and it was found that a DNA fragment of about 1.4 kb was amplified.

The obtained DNA was sequenced for the nucleotide sequences of the both ends by using the oligonucleotides of SEQ ID NOS: 1 and 2. The nucleotide sequencing was performed according to the method of Sanger (*J. Mol. Biol.*, 143, 161 (1980)) by using a DNA Sequencing Kit (produced by Applied Biosystems Co.). The determined nucleotide sequence was translated into an amino acid sequence, and compared with an amino acid sequence deduced from the *gltB* gene of *Escherichia coli* and yeast. As a result, high homology was observed. Therefore, it

was determined that the DNA fragment amplified by the PCR should be a part of the *gltB* gene of *Brevibacterium lactofermentum* ATCC13869. By using this PCR-amplified DNA fragment as a probe and a DIG DNA Labeling and  
5 Detection Kit (produced by Boehringer Mannheim), fragments obtained by digesting chromosomal DNA of *Brevibacterium lactofermentum* ATCC13869 prepared by the above method with *EcoRI*, *BamHI*, *HindIII*, *PstI* or *SalI* (produced by Takara Shuzo Co., Ltd.) were subjected to  
10 Southern hybridization in a conventional manner. AS a result, it was found that a fragment of 14 kb digested with *HindIII* was hybridized with the probe DNA.

Then, the *HindIII* fragment of chromosomal DNA of *Brevibacterium lactofermentum* ATCC13869 prepared in a  
15 conventional manner was subjected to agarose electrophoresis and a DNA fragment of about 10 kb or longer was recovered by using glass powder. The recovered DNA fragments and vector pMW219 (produced by Nippon Gene) digested with a restriction enzyme, *HindIII*  
20 (produced by Takara Shuzo Co., Ltd.), were ligated by using a ligation kit (produced by Takara Shuzo Co., Ltd.), and used for transformation of competent cells of *Escherichia coli* JM109 (produced by Takara Shuzo Co., Ltd.). The transformant strains were plated on L medium  
25 (10 g/L of Bacto trypton, 5 g/L of Bacto yeast extract, 5 g/L of NaCl, and 15 g/L of agar, pH 7.2) containing 10 µg/ml of IPTG (isopropyl-β-D-thiogalactopyranoside), 40

µg/ml of X-Gal (5-bromo-4-chloro-3-indolyl-β-D-galactoside) and 25 µg/ml of kanamycin, and cultured overnight. Then, the appeared white colonies were picked up and separated into single colonies to obtain  
5 about 1,000 transformants.

Plasmids were prepared from the obtained transformant strains by using the alkaline method (Text for Bioengineering Experiments, Edited by the Society for Bioscience and Bioengineering, Japan, p.105,  
10 Baifukan, 1992). PCR was performed under the above conditions by using as primers synthetic oligonucleotides of nucleotide sequences shown as SEQ ID NOS: 3 and 4, which were prepared based on the sequenced portion in the DNA used as a probe, and the plasmids as  
15 a template. Then, there was selected a transformant harboring a plasmid with which an amplified fragment having the same length as the DNA fragment amplified by PCR using these primers and chromosome of *Brevibacterium lactofermentum* ATCC13869 as a template, that is, about  
20 1.3 kb, could be obtained.

(2) Sequencing of DNA fragment containing *Brevibacterium lactofermentum* ATCC13869 *gltBD* gene for total nucleotide sequence and isolation of ABC transporter gene

25 The plasmid DNA prepared by the alkaline method from the transformant obtained in the above (1) contained a DNA fragment of about 14 kb derived from a

*Brevibacterium lactofermentum* ATCC13869 chromosome. The DNA fragment of about 14 kb derived from the *Brevibacterium lactofermentum* ATCC13869 chromosome in the obtained plasmid was sequenced for the total  
5 nucleotide sequence in the same manner as the method described above. As a result, it was found that, while the obtained DNA fragment contained the *gltBD* gene in the full length, it also contained two open reading frames of 500 bps or longer downstream from the *gltBD*  
10 gene in an inverted direction from the end and a sequence estimated to be a terminator downstream from these open reading frames. However, since these open reading frames lacked a promoter region, a region upstream from them was cloned as described below.

15       The region was cloned from a DNA fragment obtained through digestion of chromosome of *Brevibacterium lactofermentum* ATCC13869 with a restriction enzyme *Bam*HI by using primers of SEQ ID NOS: 5 and 6 shown in Sequence Listing and an LA PCR in vitro cloning Kit  
20 (produced by Takara Shuzo Co., Ltd.). As a result of PCR performed by using the aforementioned primers, a DNA fragment of about 1.8 kb was amplified, and hence this DNA fragment was sequenced for the nucleotide sequence in the same manner as described above. As a result, it  
25 was found that the amplified DNA fragment contained an open reading frame for about 350 amino acids located upstream from the aforementioned two open reading frames

and a region estimated to be a promoter region further upstream from it. Therefore, it is possible that these three open reading frames constitute an operon.

Nucleotide sequences of these open reading frames  
5 are shown in SEQ ID NO: 7 in Sequence Listing. Amino  
acid sequences of products deduced from the nucleotide  
sequences were also shown in SEQ ID NO: 7 in Sequence  
Listing. Among these, the nucleotide numbers 1 to 1101  
10 represent the first open reading frame, the nucleotide  
numbers 1117 to 1725 represent the second open reading  
frame and the nucleotide numbers 1759 to 2367 represent  
the third open reading frame. A methionine residue  
present at the N-terminus of the protein encoded by each  
open reading frame was derived from the initiation codon.  
15 It is well known that such a methionine residue may be  
usually irrelevant to function of the protein and  
eliminated by the action of peptidase after the  
translation. In the case of the aforementioned proteins,  
the methionine residue at the N-terminus may also be  
20 eliminated. Further, since the promoter region and  
terminator sequence estimated above were obtained just  
as a result of computerized analyses, it is possible  
that open reading frames may be present upstream or  
downstream from them and expressed together with them in  
25 fact.

The nucleotide sequences and amino acid sequences  
were compared with known sequences for homology. The

used database were EMBL and SWISS-PROT. As a result, it was found that DNA shown as SEQ ID NO: 7 in Sequence Listing and proteins encoded by it were novel genes and proteins for bacteria belonging to the genus

- 5 *Corynebacterium*. It was found that, among these, the second open reading frame and the protein encoded by it showed high homology to the already reported ATP-binding proteins of ABC transporters and the genes coding for them, and it was a gene coding for an ATP-binding
- 10 protein that was novel for bacteria belonging to the genus *Corynebacterium*.

#### Industrial Applicability

- 15 According to the present invention, constituents of ABC transporters of *Brevibacterium lactofermentum* and DNA coding for them are provided. The genes of the present invention can be utilized for breeding of coryneform bacteria.

What is claimed is:

1. A protein defined in the following (A) or (B):

(A) a protein which has the amino acid sequence of  
SEQ ID NO: 8 shown in Sequence Listing;

5 (B) a protein which has the amino acid sequence of  
SEQ ID NO: 8 shown in Sequence Listing including  
substitution, deletion, insertion, addition or inversion  
of one or several amino acids, and constitutes an ABC  
transporter.

10 2. A DNA which codes for a protein defined in the  
following (A) or (B):

(A) a protein which has the amino acid sequence of  
SEQ ID NO: 8 shown in Sequence Listing;

15 (B) a protein which has the amino acid sequence of  
SEQ ID NO: 8 shown in Sequence Listing including  
substitution, deletion, insertion, addition or inversion  
of one or several amino acids, and constitutes an ABC  
transporter.

20 3. The DNA according to Claim 2, which is a DNA  
defined in the following (a) or (b):

(a) a DNA which comprises the nucleotide sequence  
of nucleotide numbers 1 to 1101 of SEQ ID NO: 7 shown in  
Sequence Listing;

25 (b) a DNA which is hybridizable with the  
nucleotide sequence of nucleotide numbers 1 to 1101 of  
SEQ ID NO: 7 or a probe prepared from the nucleotide  
sequence under a stringent condition, and codes for a

09868338 061301

protein constituting an ABC transporter.

4. The DNA according to Claim 3, wherein the stringent condition is a condition in which washing is performed at 60°C and a salt concentration corresponding to 1 x SSC and 0.1 % SDS.

5. A protein defined in the following (C) or (D):

(C) a protein which has the amino acid sequence of SEQ ID NO: 9 shown in Sequence Listing;

(D) a protein which has the amino acid sequence of SEQ ID NO: 9 shown in Sequence Listing including substitution, deletion, insertion, addition or inversion of one or several amino acids, and has ATPase activity of ABC transporter.

6. A DNA coding for a protein defined in the following (C) or (D):

(C) a protein which has the amino acid sequence of SEQ ID NO: 9 shown in Sequence Listing;

(D) a protein which has the amino acid sequence of SEQ ID NO: 9 shown in Sequence Listing including substitution, deletion, insertion, addition or inversion of one or several amino acids, and has ATPase activity of ABC transporter.

7. The DNA according to Claim 6, which is a DNA defined in the following (c) or (d):

(c) a DNA which comprises the nucleotide sequence of nucleotide numbers 1117 to 1725 of SEQ ID NO: 7 shown in Sequence Listing;

0986333-064301



(d) a DNA which is hybridizable with the nucleotide sequence of nucleotide numbers 1117 to 1725 of SEQ ID NO: 7 or a probe prepared from the nucleotide sequence under a stringent condition, and codes for a protein having ATPase activity of ABC transporter.

8. The DNA according to Claim 7, wherein the stringent condition is a condition in which washing is performed at 60°C and a salt concentration corresponding to 1 x SSC and 0.1% SDS.

9. A protein defined in the following (E) or (F):

(E) a protein which has the amino acid sequence of SEQ ID NO: 10 shown in Sequence Listing;

(F) a protein which has the amino acid sequence of SEQ ID NO: 10 shown in Sequence Listing including substitution, deletion, insertion, addition or inversion of one or several amino acids, and constitutes an ABC transporter.

10. A DNA coding for a protein defined in the following (E) or (F):

(E) a protein which has the amino acid sequence of SEQ ID NO: 10 shown in Sequence Listing;

(F) a protein which has the amino acid sequence of SEQ ID NO: 10 shown in Sequence Listing including substitution, deletion, insertion, addition or inversion of one or several amino acids, and constitutes an ABC transporter.

11. The DNA according to Claim 10, which is a DNA

0366338 061301

defined in the following (e) or (f):

(e) a DNA which comprises the nucleotide sequence of nucleotide numbers 1759 to 2367 of SEQ ID NO: 7 shown in Sequence Listing;

5           (f) a DNA which is hybridizable with the nucleotide sequence of nucleotide numbers 1759 to 2367 of SEQ ID NO: 7 or a probe prepared from the nucleotide sequence under a stringent condition, and codes for a protein constituting an ABC transporter.

10           12. The DNA according to Claim 11, wherein the stringent condition is a condition in which washing is performed at 60°C and at a salt concentration corresponding to 1 x SSC and 0.1% SDS.

15           13. A DNA which comprises a nucleotide sequence coding for a protein having the amino acid sequence of SEQ ID NO: 8, a nucleotide sequence coding for a protein having the amino acid sequence of SEQ ID NO: 9 and a nucleotide sequence coding for a protein having the amino acid sequence of SEQ ID NO: 10.

20           14. The DNA according to Claim 13, which has the nucleotide sequence shown as SEQ ID NO: 7.

## Abstract of the Disclosure

The present invention provides constituents of ABC transporter of *Brevibacterium lactofermentum* having an amino acid sequence of SEQ ID NO: 8, 9 or 10 shown in Sequence Listing and DNAs coding for them. The DNA of the present invention can be utilized for breeding of coryneform bacteria.

09868338.061804

## Sequence Listing

<110> KANNO, Sohei  
 KIMURA, Eiichiro  
 MATSUI, Kazuhiko  
 NAKAMATSU, Tsuyoshi

<120> ABC Transporter and Gene Coding for the Same

<130> B-528SMOP924

<141> 1999-12-16

<150> JP 10-360621  
 <151> 1998-12-18

<160> 10

<170> PatentIn Ver. 2.0

<210> 1  
 <211> 17  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> UNSURE  
 <222> (3,9,12)  
 <223> n=a or c or g or t

<220>  
 <223> Description of Artificial Sequence:primer for  
 amplifying Brevibacterium lactofermentum gltBD gene

<400> 1  
 ggngarggng gngarga

<210> 2  
 <211> 18  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> UNSURE  
 <222> (1,4,7,)  
 <223> n=a or c or g or t

<220>  
 <223> Description of Artificial Sequence:primer for  
 amplifying Brevibacterium lactofermentum gltBD gene

<400> 2  
 nccncngtc atrtaytc 18

<210> 3  
 <211> 32  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:primer for  
 amplifying Brevibacterium lactofermentum gltBD gene

<400> 3  
 aatccacgtg aagctagtgg cagaacaagg cg 32

<210> 4  
 <211> 30  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:primer for  
 amplifying Brevibacterium lactofermentum gltBD gene

<400> 4  
 acgaatgaac aattcaccac tggttgcgcc 30

09863338.061804

<210> 5  
 <211> 22  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:primer for  
 amplifying downstream region of gltBD gene

<400> 5  
 atcctcgaca aggatctgtc cg 22

<210> 6  
 <211> 30  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence:primer for  
 amplifying downstream region of gltBD gene

<400> 6  
 ggtttgtaa gtgtgccaag acagttgagc 30

<210> 7  
 <211> 2370  
 <212> DNA  
 <213> Brevibacterium lactofermentum

<220>  
 <221> CDS  
 <222> (1)..(1101)

<220>  
 <221> CDS  
 <222> (1117)..(1725)

<220>  
 <221> CDS  
 <222> (1759)..(2367)

09863338.061804

atg	ctg	gcg	acc	cga	cta	att	acc	ttg	ttc	ttt	ttc	cta	gga	atc	att	48
Met	Leu	Ala	Thr	Arg	Leu	Ile	Thr	Leu	Phe	Phe	Phe	Leu	Gly	Ile	Ile	
1				5				10						15		
gga	tcg	cta	acc	ggt	aac	ctc	agt	gaa	cta	cgt	gca	caa	act	act	ttt	96
Gly	Ser	Leu	Thr	Gly	Asn	Leu	Ser	Glu	Leu	Arg	Ala	Gln	Thr	Thr	Phe	
				20				25						30		
agt	aca	tta	tgg	gat	acc	cat	aaa	gaa	acc	tat	aga	gtc	tcc	ata	gct	144
Ser	Thr	Leu	Trp	Asp	Thr	His	Lys	Glu	Thr	Tyr	Arg	Val	Ser	Ile	Ala	
				35				40						45		
tcc	gca	gca	gga	caa	gac	ttc	tac	ggg	ctt	gct	gag	act	cta	cgc	act	192
Ser	Ala	Ala	Gly	Gln	Asp	Phe	Tyr	Gly	Leu	Ala	Glu	Thr	Leu	Arg	Thr	
				50				55						60		
atg	gat	agg	cat	ggg	gaa	att	att	ttg	gca	gat	cgt	caa	tgg	tta	aca	240
Met	Asp	Arg	His	Gly	Glu	Ile	Ile	Leu	Ala	Asp	Arg	Gln	Trp	Leu	Thr	
65						70					75				80	
gct	ccc	ctt	gat	atc	ggt	gca	cca	gtc	gta	tta	tca	aac	aca	act	ttt	288
Ala	Pro	Leu	Asp	Ile	Gly	Ala	Pro	Val	Val	Leu	Ser	Asn	Thr	Thr	Phe	
				85						90					95	
gcc	gtt	gat	gaa	gga	cta	ctt	gcg	cca	aaa	gat	cta	ccg	caa	agt	gac	336
Ala	Val	Asp	Glu	Gly	Leu	Leu	Ala	Pro	Lys	Asp	Leu	Pro	Gln	Ser	Asp	
				100				105						110		
gag	atc	aca	ata	ttg	cat	cct	cag	ttt	ctg	gat	tcg	gcc	aaa	gag	cca	384
Glu	Ile	Thr	Ile	Leu	His	Pro	Gln	Phe	Leu	Asp	Ser	Ala	Lys	Glu	Pro	
				115				120						125		
gaa	tta	ctt	ggt	ttg	ctg	gag	ttc	gaa	gca	tcc	aac	tca	caa	gtg	cca	432
Glu	Leu	Leu	Gly	Leu	Leu	Glu	Phe	Glu	Ala	Ser	Asn	Ser	Gln	Val	Pro	
				130				135						140		
atg	cca	aag	atc	caa	agc	att	cca	tat	gat	agc	gaa	gac	tca	acc	aac	480
Met	Pro	Lys	Ile	Gln	Ser	Ile	Pro	Tyr	Asp	Ser	Glu	Asp	Ser	Thr	Asn	
145						150					155				160	
ccc	atg	tct	gaa	gtt	ttt	acc	tac	aac	att	aac	ctg	gat	agt	gca	gta	528
Pro	Met	Ser	Glu	Val	Phe	Thr	Tyr	Asn	Ile	Asn	Leu	Asp	Ser	Ala	Val	
				165						170					175	
aga	aac	cca	atc	gta	gtt	atc	ctt	ccc	gca	ggc	tta	gag	ctt	tta	agt	576
Arg	Asn	Pro	Ile	Val	Val	Ile	Leu	Pro	Ala	Gly	Leu	Glu	Leu	Leu	Ser	
				180						185					190	
gat	caa	aat	ttg	tcg	gct	cga	ctc	aca	cag	aat	agt	ctg	ctg	ata	aaa	624
Asp	Gln	Asn	Leu	Ser	Ala	Arg	Leu	Thr	Gln	Asn	Ser	Leu	Leu	Ile	Lys	
				195				200						205		
gac	cag	act	ggt	gtg	aac	gct	ctt	cta	tcc	tca	gag	gat	tca			

Asp Gln Thr Gly Val Asn Ala Leu Leu Ser Ser Glu Asp Ser Arg Asn	
210 215 220	
tat gtg gga gct gca tcc ccg atg att gac acg tgg gaa gaa agc gtt	720
Tyr Val Gly Ala Ala Ser Pro Met Ile Asp Thr Trp Glu Glu Ser Val	
225 230 235 240	
gtt cgg ttg aag gaa gcg aac caa ata atc gcc ttc aac gct ttc att	768
Val Arg Leu Lys Glu Ala Asn Gln Ile Ile Ala Phe Asn Ala Phe Ile	
245 250 255	
gca ttg ttc ctc acg acg act ctt gtt cta gca tac tgc act ggt att	816
Ala Leu Phe Leu Thr Thr Thr Leu Val Leu Ala Tyr Cys Thr Gly Ile	
260 265 270	
tca ttt aag aaa tca aag aag act atg ggt agc gca tct act agg aaa	864
Ser Phe Lys Lys Ser Lys Lys Thr Met Gly Ser Ala Ser Thr Arg Lys	
275 280 285	
tca tcc att aag agc tcg att aca gct gct aat tgt aga agt aat ttt	912
Ser Ser Ile Lys Ser Ser Ile Thr Ala Ala Asn Cys Arg Ser Asn Phe	
290 295 300	
cgc ttc aat tcc gtg cgt ctg gct cgc gaa ccg cta ttt cga gcg atc	960
Arg Phe Asn Ser Val Arg Leu Ala Arg Glu Pro Leu Phe Arg Ala Ile	
305 310 315 320	
tgc agc aat agc ttc aga tgc tcc ctc agc cag ata ctt aga aca tct	1008
Cys Ser Asn Ser Phe Arg Cys Ser Leu Ser Gln Ile Leu Arg Thr Ser	
325 330 335	
caa ttc tat acc tcc atc act gcc gtt ggt ttt agg aat ctt aat aat	1056
Gln Phe Tyr Thr Ser Ile Thr Ala Val Gly Phe Arg Asn Leu Asn Asn	
340 345 350	
cgg ttg gac ttc act ttc att ttt cag ttc gat gaa gct tcc ttt	1101
Arg Leu Asp Phe Thr Phe Ile Phe Gln Phe Asp Glu Ala Ser Phe	
355 360 365	
tgaaaagagc acaca atg ata gaa atc aat gac ctc aag aaa tct ttt ggc	1152
Met Ile Glu Ile Asn Asp Leu Lys Lys Ser Phe Gly	
1 5 10	
gtt cgg atc tta tgg caa ggt ctc agt cat aag ttt tta cca gga aca	1200
Val Arg Ile Leu Trp Gln Gly Leu Ser His Lys Phe Leu Pro Gly Thr	
15 20 25	
atg aca gca ctg act gga gcg tcc ggt tca gga aaa tcg act ttg ctc	1248
Met Thr Ala Leu Thr Gly Ala Ser Gly Ser Gly Lys Ser Thr Leu Leu	
30 35 40	
aac tgt ctt ggc aca ctt gac aaa cca agt tcc gga cag atc ctt gtc	1296
Asn Cys Leu Gly Thr Leu Asp Lys Pro Ser Ser Gly Gln Ile Leu Val	
45 50 55 60	

0986333 OCT 1991





60 65 70  
 gca tac ctg tgt att cat ggt gaa cct caa agc acc agc cct ttc act 2019  
 Ala Tyr Leu Cys Ile His Gly Glu Pro Gln Ser Thr Ser Pro Phe Thr  
 75 80 85  
 tta att gtt gcc caa atg gcg ttt tcg gga ttg ctc atg ttc aga ggg 2067  
 Leu Ile Val Ala Gln Met Ala Phe Ser Gly Leu Leu Met Phe Arg Gly  
 90 95 100  
 caa cgg gtg ctc gct ttt atc tct gca ggt ggg ctc att tgg att ggg 2115  
 Gln Arg Val Leu Ala Phe Ile Ser Ala Gly Gly Leu Ile Trp Ile Gly  
 105 110 115  
 acc atc gat cca aca aac ggt gct tgg tct cct cat gtg atg tcc gcg 2163  
 Thr Ile Asp Pro Thr Asn Gly Ala Trp Ser Pro His Val Met Ser Ala  
 120 125 130 135  
 cta gca ctt gcc gta ttc ttt gcg ctg tcg atg gca ctt gga cag gtt 2211  
 Leu Ala Leu Ala Val Phe Phe Ala Leu Ser Met Ala Leu Gly Gln Val  
 140 145 150  
 ctt cga tca aaa gtt gaa caa aga gcc aac ctt gag gag cag gca aaa 2259  
 Leu Arg Ser Lys Val Glu Gln Arg Ala Asn Leu Glu Glu Gln Ala Lys  
 155 160 165  
 att cag aca gaa ctg cgc aga aaa gaa cta agc act cca tct gca tcg 2307  
 Ile Gln Thr Glu Leu Arg Arg Lys Glu Leu Ser Thr Pro Ser Ala Ser  
 170 175 180  
 gtc ggt tgc caa aga act tac gtt tgc agt gat gaa atc gca gga gct 2355  
 Val Gly Cys Gln Arg Thr Tyr Val Cys Ser Asp Glu Ile Ala Gly Ala  
 185 190 195  
 cag tgg tcg cga taa 2370  
 Gln Trp Ser Arg  
 200

&lt;210&gt; 8

&lt;211&gt; 367

&lt;212&gt; PRT

<213> *Brevibacterium lactofermentum*

&lt;400&gt; 8

Met Leu Ala Thr Arg Leu Ile Thr Leu Phe Phe Phe Leu Gly Ile Ile  
 1 5 10 15  
 Gly Ser Leu Thr Gly Asn Leu Ser Glu Leu Arg Ala Gln Thr Thr Phe  
 20 25 30  
 Ser Thr Leu Trp Asp Thr His Lys Glu Thr Tyr Arg Val Ser Ile Ala  
 35 40 45

09868338.061801

Ser Ala Ala Gly Gln Asp Phe Tyr Gly Leu Ala Glu Thr Leu Arg Thr  
 50 55 60  
 Met Asp Arg His Gly Glu Ile Ile Leu Ala Asp Arg Gln Trp Leu Thr  
 65 70 75 80  
 Ala Pro Leu Asp Ile Gly Ala Pro Val Val Leu Ser Asn Thr Thr Phe  
 85 90 95  
 Ala Val Asp Glu Gly Leu Leu Ala Pro Lys Asp Leu Pro Gln Ser Asp  
 100 105 110  
 Glu Ile Thr Ile Leu His Pro Gln Phe Leu Asp Ser Ala Lys Glu Pro  
 115 120 125  
 Glu Leu Leu Gly Leu Leu Glu Phe Glu Ala Ser Asn Ser Gln Val Pro  
 130 135 140  
 Met Pro Lys Ile Gln Ser Ile Pro Tyr Asp Ser Glu Asp Ser Thr Asn  
 145 150 155 160  
 Pro Met Ser Glu Val Phe Thr Tyr Asn Ile Asn Leu Asp Ser Ala Val  
 165 170 175  
 Arg Asn Pro Ile Val Val Ile Leu Pro Ala Gly Leu Glu Leu Leu Ser  
 180 185 190  
 Asp Gln Asn Leu Ser Ala Arg Leu Thr Gln Asn Ser Leu Leu Ile Lys  
 195 200 205  
 Asp Gln Thr Gly Val Asn Ala Leu Leu Ser Ser Glu Asp Ser Arg Asn  
 210 215 220  
 Tyr Val Gly Ala Ala Ser Pro Met Ile Asp Thr Trp Glu Glu Ser Val  
 225 230 235 240  
 Val Arg Leu Lys Glu Ala Asn Gln Ile Ile Ala Phe Asn Ala Phe Ile  
 245 250 255  
 Ala Leu Phe Leu Thr Thr Thr Leu Val Leu Ala Tyr Cys Thr Gly Ile  
 260 265 270  
 Ser Phe Lys Lys Ser Lys Lys Thr Met Gly Ser Ala Ser Thr Arg Lys  
 275 280 285  
 Ser Ser Ile Lys Ser Ser Ile Thr Ala Ala Asn Cys Arg Ser Asn Phe  
 290 295 300  
 Arg Phe Asn Ser Val Arg Leu Ala Arg Glu Pro Leu Phe Arg Ala Ile  
 305 310 315 320  
 Cys Ser Asn Ser Phe Arg Cys Ser Leu Ser Gln Ile Leu Arg Thr Ser  
 325 330 335  
 Gln Phe Tyr Thr Ser Ile Thr Ala Val Gly Phe Arg Asn Leu Asn Asn  
 340 345 350  
 Arg Leu Asp Phe Thr Phe Ile Phe Gln Phe Asp Glu Ala Ser Phe  
 355 360 365

09868338 061801

<210> 9  
 <211> 203  
 <212> PRT  
 <213> Brevibacterium lactofermentum

<400> 9

Met Ile Glu Ile Asn Asp Leu Lys Lys Ser Phe Gly Val Arg Ile Leu  
 1 5 10 15  
 Trp Gln Gly Leu Ser His Lys Phe Leu Pro Gly Thr Met Thr Ala Leu  
 20 25 30  
 Thr Gly Ala Ser Gly Ser Gly Lys Ser Thr Leu Leu Asn Cys Leu Gly  
 35 40 45  
 Thr Leu Asp Lys Pro Ser Ser Gly Gln Ile Leu Val Glu Asp Val Asp  
 50 55 60  
 Leu Leu Lys Leu Ser Thr Arg Lys Gln Arg Leu Tyr Arg Lys Asn Thr  
 65 70 75 80  
 Val Gly Tyr Leu Phe Gln Asp Tyr Ala Leu Ile Pro Asp Arg Thr Val  
 85 90 95  
 Lys Phe Asn Leu Gln Leu Ala Val Glu Lys His Lys Trp Pro Glu Ile  
 100 105 110  
 Pro Gln Val Leu His Ala Val Gly Leu Glu Ser Phe Glu Glu Lys Pro  
 115 120 125  
 Val Phe Glu Leu Ser Gly Gly Glu Gln Gln Arg Thr Ala Leu Ala Arg  
 130 135 140  
 Val Leu Leu Lys Asn Pro Arg Ile Ile Leu Ala Asp Glu Pro Thr Gly  
 145 150 155 160  
 Ala Leu Asp Leu Thr Asn Ser Glu Leu Val Ile Glu Ala Leu Arg Ala  
 165 170 175  
 Leu Ala Asp Lys Gly Ala Thr Val Val Ala Thr His Ser Pro Leu  
 180 185 190  
 Phe Arg Glu Ser Ala Asp Thr Ile Ile Lys Leu  
 195 200

<210> 10  
 <211> 203  
 <212> PRT  
 <213> Brevibacterium lactofermentum

<400> 10

Met Met Glu Phe Leu Asn Thr His Arg Leu Ile Val Leu Gly Ser Leu  
 1 5 10 15

09060333 061801

Ser Phe Leu Gly Leu Gly Phe Ala Glu Val Leu Leu Arg Gly Gln Trp  
           20                          25                          30  
 Ser Thr Pro Gln Phe Phe Val Phe Thr Phe Leu Gln Thr Leu Leu Leu  
           35                          40                          45  
 Val Leu Cys Phe Ile Pro Lys Leu Ser Val Pro Phe Val Val Leu Leu  
           50                          55                          60  
 Ser Ile Ala Gln Leu Ala Leu Ala Tyr Leu Cys Ile His Gly Glu Pro  
           65                          70                          75                          80  
 Gln Ser Thr Ser Pro Phe Thr Leu Ile Val Ala Gln Met Ala Phe Ser  
                           85                          90                          95  
 Gly Leu Leu Met Phe Arg Gly Gln Arg Val Leu Ala Phe Ile Ser Ala  
                           100                          105                          110  
 Gly Gly Leu Ile Trp Ile Gly Thr Ile Asp Pro Thr Asn Gly Ala Trp  
                           115                          120                          125  
 Ser Pro His Val Met Ser Ala Leu Ala Leu Ala Val Phe Phe Ala Leu  
           130                          135                          140  
 Ser Met Ala Leu Gly Gln Val Leu Arg Ser Lys Val Glu Gln Arg Ala  
           145                          150                          155                          160  
 Asn Leu Glu Glu Gln Ala Lys Ile Gln Thr Glu Leu Arg Arg Lys Glu  
                           165                          170                          175  
 Leu Ser Thr Pro Ser Ala Ser Val Gly Cys Gln Arg Thr Tyr Val Cys  
                           180                          185                          190  
 Ser Asp Glu Ile Ala Gly Ala Gln Trp Ser Arg  
           195                          200

09868338.061801

# Declaration, Power Of Attorney and Petition

Page 1 of 3

WE (I) the undersigned inventor(s), hereby declare(s) that:

My residence, post office address and citizenship are as stated below next to my name,

We (I) believe that we are (I am) the original, first, and joint (sole) inventor(s) of the subject matter which is claimed and for which a patent is sought on the invention entitled

ABC TRANSPORTER AND GENE CODING FOR THE SAME

the specification of which

☒ is attached hereto.

☐ was filed on \_\_\_\_\_ as

Application Serial No. \_\_\_\_\_

and amended on \_\_\_\_\_.

☒ was filed as PCT international application

Number PCT/JP 99/07079

on December 16, 1999,

and was amended under PCT Article 19

on \_\_\_\_\_ (if applicable).

We (I) hereby state that we (I) have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

We (I) acknowledge the duty to disclose information known to be material to the patentability of this application as defined in Section 1.56 of Title 37 Code of Federal Regulations.

We (I) hereby claim foreign priority benefits under 35 U.S.C. § 119(a)-(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate, or § 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or PCT International application having a filing date before that of the application on which priority is claimed. Prior Foreign Application(s)

Application No.	Country	Day/Month/Year	Priority Claimed
10-360621	Japan	18/12/1998	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No
_____	_____	_____	<input type="checkbox"/> Yes <input type="checkbox"/> No
_____	_____	_____	<input type="checkbox"/> Yes <input type="checkbox"/> No
_____	_____	_____	<input type="checkbox"/> Yes <input type="checkbox"/> No

We (I) hereby claim the benefit under Title 35, United States Code, § 119(e) of any United States provisional application(s) listed below.

_____	_____
(Application Number)	(Filing Date)
_____	_____
(Application Number)	(Filing Date)

We (I) hereby claim the benefit under 35 U.S.C. § 120 of any United States application(s), or § 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. § 112, I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR § 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application.

Application Serial No.	Filing Date	Status (pending, patented, abandoned)
_____	_____	_____
_____	_____	_____
_____	_____	_____

And we (I) hereby appoint: Norman F. Oblon, Registration Number 24,618; Marvin J. Spivak, Registration Number 24,913; C. Irvin McClelland, Registration Number 21,124; Gregory J. Maier, Registration Number 25,599; Arthur I. Neustadt, Registration Number 24,854; Richard D. Kelly, Registration Number 27,757; James D. Hamilton, Registration Number 28,421; Eckhard H. Kuesters, Registration Number 28,870; Robert T. Pous, Registration Number 29,099; Charles L. Gholz, Registration Number 26,395; Vincent J. Sunderdick, Registration Number 29,004; William E. Beaumont, Registration Number 30,996; Steven B. Kelber, Registration Number 30,073; Robert F. Gnuse, Registration Number 27,295; Jean-Paul Lavallee, Registration Number 31,451; Timothy R. Schwartz, Registration Number 32,171; Stephen G. Baxter, Registration Number 32,884; Martin M. Zoltick, Registration Number 35,745; Robert W. Hahl, Registration Number 33,893; Richard L. Treanor, Registration Number 36,379; Steven P. Weihrouch, Registration Number 32,829; John T. Goolkasian, Registration Number 26,142; Marc R. Labgold, Registration Number 34,651; William J. Healey, Registration Number 36,160; and Richard L. Chinn, Registration Number 34,305; our (my) attorneys, with full powers of substitution and revocation, to prosecute this application and to transact all business in the Patent Office connected therewith; and we (I) hereby request that all correspondence regarding this application be sent to the firm of OBLON, SPIVAK, McCLELLAND, MAIER & NEUSTADT, P.C., whose Post Office Address is: Fourth Floor, 1755 Jefferson Davis Highway, Arlington, Virginia 22202.

We (I) declare that all statements made herein of our (my) own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Sohei KANNNO

NAME OF FIRST SOLE INVENTOR

Residence: Kawasaki-shi, Kanagawa, Japan

JPX

Sohei Kanno

Signature of Inventor

Citizen of: Japan

Post Office Address: c/o Ajinomoto Co., Inc.,  
Fermentation & Biotechnology Laboratories,  
1-1, Suzuki-cho, Kawasaki-ku, Kawasaki-shi,  
Kanagawa, Japan

June 5, 2001

Date

Eiichiro KIMURA

NAME OF SECOND JOINT INVENTOR

*Eiichiro Kimura*

Signature of Inventor

June 5, 2001

Date

Residence: Kawasaki-shi, Kanagawa, Japan

JPX

Citizen of: Japan

Post Office Address: c/o Ajinomoto Co., Inc.,  
Fermentation & Biotechnology Laboratories,  
1-1, Suzuki-cho, Kawasaki-ku, Kawasaki-shi,  
Kanagawa, Japan

Kazuhiko MATSUI

NAME OF THIRD JOINT INVENTOR

*Kazuhiko Matsui*

Signature of Inventor

June 5, 2001

Date

Residence: Kawasaki-shi, Kanagawa, Japan

JPX

Citizen of: Japan

Post Office Address: c/o Ajinomoto Co., Inc.,  
Fermentation & Biotechnology Laboratories,  
1-1, Suzuki-cho, Kawasaki-ku, Kawasaki-shi,  
Kanagawa, Japan

Tsuyoshi NAKAMATSU

NAME OF FOURTH JOINT INVENTOR

*Tsuyoshi Nakamatsu*

Signature of Inventor

June 5, 2001

Date

Residence: Kawasaki-shi, Kanagawa, Japan

JPX

Citizen of: Japan

Post Office Address: c/o Ajinomoto Co., Inc.,  
Fermentation & Biotechnology Laboratories,  
1-1, Suzuki-cho, Kawasaki-ku, Kawasaki-shi,  
Kanagawa, Japan

NAME OF FIFTH JOINT INVENTOR

Signature of Inventor

Date



SEQUENCE LISTING

<110> KANNO , SOHEI  
MATSUI, KAZUHIKO  
NAKAMATSU, TSUYOSHI  
KIMURA, EIICHIRO

<120> ABC TRANSPORTER AND GENE CODING FOR THE SAME

<130> 209861US-8222-10-0-PCT

<140> US09/868338

<141> 2001-06-18

<150> JP 10/360621

<151> 1998-12-18

<150> PCT/JP 99/07079

<151> 1998-12-18

<160> 10

<170> PatentIn version 3.1

<210> 1

<211> 17

<212> DNA

<213> ARTIFICIAL SEQUENCE

<220>

<223> SYNTHETIC DNA

<220>

<221> misc\_feature

<222> (3)..(3)

<223> n = c, g, a, or t

<220>

<221> misc\_feature

<222> (9)..(9)

<223> n = c, g, a, or t

<220>

<221> misc\_feature

<222> (12)..(12)

<223> n = c, g, a, or t

<400> 1

ggngarggng gngarga

17

<210> 2

<211> 18

<212> DNA

<213> ARTIFICIAL SEQUENCE

<220>

<223> SYNTHETIC DNA

<220>

<221> misc\_feature

<222> (1)..(1)

<223> n = c, g, a, or t

<220>

<221> misc\_feature

<222> (4)..(4)

<223> n = c, g, a, or t

<220>

<221> misc\_feature

<222> (7)..(7)

<223> n = c, g, a, or t

<400> 2

nccnccngtc atrtaytc

18

<210> 3

<211> 32

<212> DNA

<213> ARTIFICIAL SEQUENCE

<220>

<223> SYNTHETIC DNA

<400> 3

aatccacgtg aagctagtgg cagaacaagg cg

32

<210> 4

<211> 30

<212> DNA

<213> ARTIFICIAL SEQUENCE

<220>

<223> SYNTHETIC DNA

<400> 4

acgaatgaac aattcaccac tggttgcgcc

30

<210> 5

<211> 22

<212> DNA

<213> ARTIFICIAL SEQUENCE

<220>

<223> SYNTHETIC DNA

<400> 5

atcctcgaca aggatctgtc cg

22

<210> 6

<211> 30

<212> DNA

<213> ARTIFICIAL SEQUENCE

<220>

<223> SYNTHETIC DNA

<400> 6

ggtttgtaa gtgtgccaag acagttgagc

30

<210> 7



gag Glu	atc Ile	aca Thr 115	ata Ile	ttg Leu	cat His	cct Pro	cag Gln 120	ttt Phe	ctg Leu	gat Asp	tcg Ser	gcc Ala 125	aaa Lys	gag Glu	cca Pro	384
gaa Glu 130	tta Leu	ctt Leu	ggt Gly	ttg Leu	ctg Leu	gag Glu 135	ttc Phe	gaa Glu	gca Ala	tcc Ser	aac Asn 140	tca Ser	caa Gln	gtg Val	cca Pro	432
atg Met 145	cca Pro	aag Lys	atc Ile	caa Gln	agc Ser 150	att Ile	cca Pro	tat Tyr	gat Asp	agc Ser 155	gaa Glu	gac Asp	tca Ser	acc Thr	aac Asn 160	480
ccc Pro	atg Met	tct Ser	gaa Glu	gtt Val 165	ttt Phe	acc Thr	tac Tyr	aac Asn 170	att Ile	aac Asn	ctg Leu	gat Asp	agt Ser	gca Ala 175	gta Val	528
aga Arg	aac Asn	cca Pro	atc Ile 180	gta Val	gtt Val	atc Ile	ctt Leu 185	ccc Pro	gca Ala	ggc Gly	tta Leu 190	gag Glu	ctt Leu	tta Leu	agt Ser	576
gat Asp	caa Gln 195	aat Asn	ttg Leu	tcg Ser	gct Ala	cga Arg 200	ctc Leu	aca Thr	cag Gln	aat Asn	agt Ser 205	ctg Leu	ctg Leu	ata Ile	aaa Lys	624
gac Asp 210	cag Gln	act Thr	ggt Gly	gtg Val	aac Asn 215	gct Ala	ctt Leu	cta Leu	tcc Ser	tca Ser 220	gag Glu	gat Asp	tca Ser	cgc Arg	aat Asn	672
tat Tyr 225	gtg Val	gga Gly	gct Ala	gca Ala	tcc Ser 230	ccg Pro	atg Met	att Ile	gac Asp 235	acg Thr	tgg Trp	gaa Glu	gaa Glu	agc Ser	gtt Val 240	720
gtt Val	cgg Arg	ttg Leu	aag Lys 245	gaa Glu	gcg Ala	aac Asn	caa Gln	ata Ile 250	atc Ile	gcc Ala	ttc Phe	aac Asn	gct Ala	ttc Phe 255	att Ile	768
gca Ala	ttg Leu	ttc Phe	ctc Leu 260	acg Thr	acg Thr	act Thr	ctt Leu 265	gtt Val	cta Leu	gca Ala	tac Tyr	tgc Cys 270	act Thr	ggt Gly	att Ile	816
tca Ser	ttt Phe 275	aag Lys	aaa Lys	tca Ser	aag Lys	aag Lys	act Thr 280	atg Met	ggg Gly	agc Ser	gca Ala 285	tct Ser	act Thr	agg Arg	aaa Lys	864
tca Ser 290	tcc Ser	att Ile	aag Lys	agc Ser	tcg Ser	att Ile 295	aca Thr	gct Ala	gct Ala	aat Asn 300	tgt Cys	aga Arg	agt Ser	aat Asn	ttt Phe	912
cgc Arg 305	ttc Phe	aat Asn	tcc Ser	gtg Val	cgt Arg 310	ctg Leu	gct Ala	cgc Arg	gaa Glu	ccg Pro 315	cta Leu	ttt Phe	cga Arg	gcg Ala	atc Ile 320	960
tgc Cys	agc Ser	aat Asn	agc Ser	ttc Phe 325	aga Arg	tgc Cys	tcc Ser	ctc Leu 330	agc Ser	cag Gln	ata Ile	ctt Leu	aga Arg	aca Thr 335	tct Ser	1008

caa	ttc	tat	acc	tcc	atc	act	gcc	gtt	ggt	ttt	agg	aat	ctt	aat	aat	1056
Gln	Phe	Tyr	Thr	Ser	Ile	Thr	Ala	Val	Gly	Phe	Arg	Asn	Leu	Asn	Asn	
			340					345					350			
cgg	ttg	gac	ttc	act	ttc	att	ttt	cag	ttc	gat	gaa	gct	tcc	ttt		1101
Arg	Leu	Asp	Phe	Thr	Phe	Ile	Phe	Gln	Phe	Asp	Glu	Ala	Ser	Phe		
		355					360					365				
tgaaaagagc	acaca	atg	ata	gaa	atc	aat	gac	ctc	aag	aaa	tct	ttt	ggc			1152
		Met	Ile	Glu	Ile	Asn	Asp	Leu	Lys	Lys	Ser	Phe	Gly			
				370						375						
gtt	cgg	atc	tta	tgg	caa	ggt	ctc	agt	cat	aag	ttt	tta	cca	gga	aca	1200
Val	Arg	Ile	Leu	Trp	Gln	Gly	Leu	Ser	His	Lys	Phe	Leu	Pro	Gly	Thr	
380					385					390					395	
atg	aca	gca	ctg	act	gga	gcg	tcc	ggt	tca	gga	aaa	tcg	act	ttg	ctc	1248
Met	Thr	Ala	Leu	Thr	Gly	Ala	Ser	Gly	Ser	Gly	Lys	Ser	Thr	Leu	Leu	
			400					405						410		
aac	tgt	ctt	ggc	aca	ctt	gac	aaa	cca	agt	tcc	gga	cag	atc	ctt	gtc	1296
Asn	Cys	Leu	Gly	Thr	Leu	Asp	Lys	Pro	Ser	Ser	Gly	Gln	Ile	Leu	Val	
		415					420						425			
gag	gat	gta	gac	ctt	ctg	aaa	ctc	tct	acg	cgt	aag	caa	cgg	tta	tac	1344
Glu	Asp	Val	Asp	Leu	Leu	Lys	Leu	Ser	Thr	Arg	Lys	Gln	Arg	Leu	Tyr	
		430					435					440				
agg	aaa	aat	acg	gtg	ggc	tat	tta	ttt	caa	gat	tat	gcc	ttg	att	ccc	1392
Arg	Lys	Asn	Thr	Val	Gly	Tyr	Leu	Phe	Gln	Asp	Tyr	Ala	Leu	Ile	Pro	
	445					450					455					
gac	agg	aca	gtt	aaa	ttc	aac	ctt	cag	ctt	gcg	gtg	gaa	aaa	cac	aaa	1440
Asp	Arg	Thr	Val	Lys	Phe	Asn	Leu	Gln	Leu	Ala	Val	Glu	Lys	His	Lys	
460				465						470					475	
tgg	cct	gaa	att	cct	caa	gta	ctt	cat	gct	gtt	ggg	ctt	gag	tcg	ttc	1488
Trp	Pro	Glu	Ile	Pro	Gln	Val	Leu	His	Ala	Val	Gly	Leu	Glu	Ser	Phe	
			480						485					490		
gag	gaa	aag	cca	gtt	ttt	gaa	ctc	tct	ggt	ggc	gaa	caa	caa	cga	act	1536
Glu	Glu	Lys	Pro	Val	Phe	Glu	Leu	Ser	Gly	Gly	Glu	Gln	Gln	Arg	Thr	
			495					500					505			
gcg	ttg	gcc	cgg	gta	ctg	ctc	aaa	aat	ccc	cga	ata	att	ctg	gct	gat	1584
Ala	Leu	Ala	Arg	Val	Leu	Leu	Lys	Asn	Pro	Arg	Ile	Ile	Leu	Ala	Asp	
		510					515					520				
gaa	cca	acc	gga	gct	cta	gat	tta	aca	aac	agt	gag	cta	gtc	ata	gaa	1632
Glu	Pro	Thr	Gly	Ala	Leu	Asp	Leu	Thr	Asn	Ser	Glu	Leu	Val	Ile	Glu	
	525					530					535					
gca	ttg	aga	gca	ctc	gcc	gac	aaa	ggc	gcc	acc	gtt					

His Ser Pro Leu Phe Arg Glu Ser Ala Asp Thr Ile Ile Lys Leu	
560 565 570	
taggtgcccc aacttttcgag agatctcagtg gca atg atg gaa ttc tta aac act	1779
Met Met Glu Phe Leu Asn Thr	575
cac cgt ttg att gtt ctc ggg agt ttg tct ttt cta ggg cta ggt ttc	1827
His Arg Leu Ile Val Leu Gly Ser Leu Ser Phe Leu Gly Leu Gly Phe	580 585 590
gcg gaa gtc ctg ctg cgt ggc cag tgg tca aca ccg cag ttt ttt gtt	1875
Ala Glu Val Leu Leu Arg Gly Gln Trp Ser Thr Pro Gln Phe Phe Val	595 600 605
ttc act ttc ttg caa act ctg ctt ctc gta ttg tgt ttt att cct aaa	1923
Phe Thr Phe Leu Gln Thr Leu Leu Leu Val Leu Cys Phe Ile Pro Lys	610 615 620 625
ctc tcg gtt cct ttt gtg gtg ctt cta agc att gcc caa ctc gcg ctt	1971
Leu Ser Val Pro Phe Val Val Leu Leu Ser Ile Ala Gln Leu Ala Leu	630 635 640
gca tac ctg tgt att cat ggt gaa cct caa agc acc agc cct ttc act	2019
Ala Tyr Leu Cys Ile His Gly Glu Pro Gln Ser Thr Ser Pro Phe Thr	645 650 655
tta att gtt gcc caa atg gcg ttt tcg gga ttg ctc atg ttc aga ggg	2067
Leu Ile Val Ala Gln Met Ala Phe Ser Gly Leu Leu Met Phe Arg Gly	660 665 670
caa cggtgtg ctc gct ttt atc tct gca ggt ggg ctc att tgg att ggg	2115
Gln Arg Val Leu Ala Phe Ile Ser Ala Gly Gly Leu Ile Trp Ile Gly	675 680 685
acc atc gat cca aca aac ggt gct tgg tct cct cat gtg atg tcc gcg	2163
Thr Ile Asp Pro Thr Asn Gly Ala Trp Ser Pro His Val Met Ser Ala	690 695 700 705
cta gca ctt gcc gta ttc ttt gcg ctg tcg atg gca ctt gga cag gtt	2211
Leu Ala Leu Ala Val Phe Phe Ala Leu Ser Met Ala Leu Gly Gln Val	710 715 720
ctt cga tca aaa gtt gaa caa aga gcc aac ctt gag gag cag gca aaa	2259
Leu Arg Ser Lys Val Glu Gln Arg Ala Asn Leu Glu Glu Gln Ala Lys	725 730 735
att cag aca gaa ctg cgc aga aaa gaa cta agc act cca tct gca tcg	2307
Ile Gln Thr Glu Leu Arg Arg Lys Glu Leu Ser Thr Pro Ser Ala Ser	740 745 750
gtc ggt tgc caa aga act tac gtt tgc agt gat gaa atc gca gga gct	2355
Val Gly Cys Gln Arg Thr Tyr Val Cys Ser Asp Glu Ile Ala Gly Ala	755 760 765
cag tgg tcg cga taa	2370
Gln Trp Ser Arg	

770

<210> 8

<211> 367

<212> PRT

<213> Brevibacterium lactofermentum

<400> 8

Met Leu Ala Thr Arg Leu Ile Thr Leu Phe Phe Phe Leu Gly Ile Ile  
1 5 10 15

Gly Ser Leu Thr Gly Asn Leu Ser Glu Leu Arg Ala Gln Thr Thr Phe  
20 25 30

Ser Thr Leu Trp Asp Thr His Lys Glu Thr Tyr Arg Val Ser Ile Ala  
35 40 45

Ser Ala Ala Gly Gln Asp Phe Tyr Gly Leu Ala Glu Thr Leu Arg Thr  
50 55 60

Met Asp Arg His Gly Glu Ile Ile Leu Ala Asp Arg Gln Trp Leu Thr  
65 70 75 80

Ala Pro Leu Asp Ile Gly Ala Pro Val Val Leu Ser Asn Thr Thr Phe  
85 90 95

Ala Val Asp Glu Gly Leu Leu Ala Pro Lys Asp Leu Pro Gln Ser Asp  
100 105 110

Glu Ile Thr Ile Leu His Pro Gln Phe Leu Asp Ser Ala Lys Glu Pro  
115 120 125

Glu Leu Leu Gly Leu Leu Glu Phe Glu Ala Ser Asn Ser Gln Val Pro  
130 135 140

Met Pro Lys Ile Gln Ser Ile Pro Tyr Asp Ser Glu Asp Ser Thr Asn  
145 150 155 160

Pro Met Ser Glu Val Phe Thr Tyr Asn Ile Asn Leu Asp Ser Ala Val  
165 170 175

Arg Asn Pro Ile Val Val Ile Leu Pro Ala Gly Leu Glu Leu Leu Ser



09063338 061304

180

185

190

Asp Gln Asn Leu Ser Ala Arg Leu Thr Gln Asn Ser Leu Leu Ile Lys  
195 200 205

Asp Gln Thr Gly Val Asn Ala Leu Leu Ser Ser Glu Asp Ser Arg Asn  
210 215 220

Tyr Val Gly Ala Ala Ser Pro Met Ile Asp Thr Trp Glu Glu Ser Val  
225 230 235 240

Val Arg Leu Lys Glu Ala Asn Gln Ile Ile Ala Phe Asn Ala Phe Ile  
245 250 255

Ala Leu Phe Leu Thr Thr Thr Leu Val Leu Ala Tyr Cys Thr Gly Ile  
260 265 270

Ser Phe Lys Lys Ser Lys Lys Thr Met Gly Ser Ala Ser Thr Arg Lys  
275 280 285

Ser Ser Ile Lys Ser Ser Ile Thr Ala Ala Asn Cys Arg Ser Asn Phe  
290 295 300

Arg Phe Asn Ser Val Arg Leu Ala Arg Glu Pro Leu Phe Arg Ala Ile  
305 310 315 320

Cys Ser Asn Ser Phe Arg Cys Ser Leu Ser Gln Ile Leu Arg Thr Ser  
325 330 335

Gln Phe Tyr Thr Ser Ile Thr Ala Val Gly Phe Arg Asn Leu Asn Asn  
340 345 350

Arg Leu Asp Phe Thr Phe Ile Phe Gln Phe Asp Glu Ala Ser Phe  
355 360 365

<210> 9

<211> 203

<212> PRT

<213> Brevibacterium lactofermentum

<400> 9

Met Ile Glu Ile Asn Asp Leu Lys Lys Ser Phe Gly Val Arg Ile Leu

1                      5                      10                      15  
 Trp Gln Gly Leu Ser His Lys Phe Leu Pro Gly Thr Met Thr Ala Leu  
                     20                      25                      30  
 Thr Gly Ala Ser Gly Ser Gly Lys Ser Thr Leu Leu Asn Cys Leu Gly  
                     35                      40                      45  
 Thr Leu Asp Lys Pro Ser Ser Gly Gln Ile Leu Val Glu Asp Val Asp  
                     50                      55                      60  
 Leu Leu Lys Leu Ser Thr Arg Lys Gln Arg Leu Tyr Arg Lys Asn Thr  
                     65                      70                      75                      80  
 Val Gly Tyr Leu Phe Gln Asp Tyr Ala Leu Ile Pro Asp Arg Thr Val  
                     85                      90                      95  
 Lys Phe Asn Leu Gln Leu Ala Val Glu Lys His Lys Trp Pro Glu Ile  
                     100                      105                      110  
 Pro Gln Val Leu His Ala Val Gly Leu Glu Ser Phe Glu Glu Lys Pro  
                     115                      120                      125  
 Val Phe Glu Leu Ser Gly Gly Glu Gln Gln Arg Thr Ala Leu Ala Arg  
                     130                      135                      140  
 Val Leu Leu Lys Asn Pro Arg Ile Ile Leu Ala Asp Glu Pro Thr Gly  
                     145                      150                      155                      160  
 Ala Leu Asp Leu Thr Asn Ser Glu Leu Val Ile Glu Ala Leu Arg Ala  
                     165                      170                      175  
 Leu Ala Asp Lys Gly Ala Thr Val Val Val Ala Thr His Ser Pro Leu  
                     180                      185                      190  
 Phe Arg Glu Ser Ala Asp Thr Ile Ile Lys Leu  
                     195                      200

<210> 10

<211> 203

<212> PRT

<213> Brevibacterium lactofermentum

<400> 10

Met Met Glu Phe Leu Asn Thr His Arg Leu Ile Val Leu Gly Ser Leu  
1 5 10 15

Ser Phe Leu Gly Leu Gly Phe Ala Glu Val Leu Leu Arg Gly Gln Trp  
20 25 30

Ser Thr Pro Gln Phe Phe Val Phe Thr Phe Leu Gln Thr Leu Leu Leu  
35 40 45

Val Leu Cys Phe Ile Pro Lys Leu Ser Val Pro Phe Val Val Leu Leu  
50 55 60

Ser Ile Ala Gln Leu Ala Leu Ala Tyr Leu Cys Ile His Gly Glu Pro  
65 70 75 80

Gln Ser Thr Ser Pro Phe Thr Leu Ile Val Ala Gln Met Ala Phe Ser  
85 90 95

Gly Leu Leu Met Phe Arg Gly Gln Arg Val Leu Ala Phe Ile Ser Ala  
100 105 110

Gly Gly Leu Ile Trp Ile Gly Thr Ile Asp Pro Thr Asn Gly Ala Trp  
115 120 125

Ser Pro His Val Met Ser Ala Leu Ala Leu Ala Val Phe Phe Ala Leu  
130 135 140

Ser Met Ala Leu Gly Gln Val Leu Arg Ser Lys Val Glu Gln Arg Ala  
145 150 155 160

Asn Leu Glu Glu Gln Ala Lys Ile Gln Thr Glu Leu Arg Arg Lys Glu  
165 170 175

Leu Ser Thr Pro Ser Ala Ser Val Gly Cys Gln Arg Thr Tyr Val Cys  
180 185 190

Ser Asp Glu Ile Ala Gly Ala Gln Trp Ser Arg  
195 200